



Phragmites!



©2002 Gary Fewless



Reed Canary



White Sweet Clover



Thistle & Yellow Clover

Month	Visits by Crew	Herbicide	Brush Clearing	Mow	Collect & Disperse	Burn Prep	Burn **
April	0-1	X	X			X	
May	1-2	X	X	X	X	X	
June	1-2	X		X	X		
July	1-2	X		X			
August	1-2	X		X			
September	0-1	X		X			
October	1-2	X	X	X	X	X	
November	0-1	X	X		X	X	
December through March	0-1	X	X		X	X	

Glyphosate (Trade Name: Razor Pro)

Category: Non-Selective

Target Species: Herbaceous & Woody – Upland / Terrestrial Areas

Application Methods: Foliar Broadcast Spray, Foliar Wick, Foliar Wiper, Cut Stem (Sponge or Low Pressure Sprayer)

Note: Needs to be above freezing, rain-fast in one hour.

Otherwise, used year-round

Aquatic-Approved Glyphosate (Trade Name: Aquaneat)

Category: Non-Selective

Target Species: Herbaceous & Woody – Wetlands / Aquatic Sites

Application Methods: Foliar Broadcast Spray, Foliar Wick, Foliar Wiper, Cut Stem (Sponge or Low Pressure Sprayer)

Note: Needs to be above freezing, rain-fast in one hour

Otherwise, used year-round

Clopyralid (Trade Name: Transline)

Category: Partially Selective

Target Species: Herbaceous & Woody Legumes, Herbaceous Composites –
Upland / Terrestrial Areas

Application Methods: Foliar Broadcast Spray, Foliar Wick, Foliar Wiper, Cut
Stem (Sponge or Low Pressure Sprayer)

Note: Best used when temps are below 79 degrees F or above 25 degrees F,
Rain-fast in two hours

2,4-D Amine (Trade Name: Weedar 64)

Category: Selective

Target Species: Herbaceous Broadleaves – Upland / Terrestrial Areas

Application Methods: Foliar Broadcast Spray

Note: Best used when temps are below 84 degrees F, Rain-fast in four hours



Chemical Cocktail

Triclopyr Amine (Trade Name: Tahoe 3a)

Category: Selective

Target Species: Herbaceous Broadleaves – Wetlands / Aquatic Sites

Application Methods: Foliar Broadcast Spray, Foliar Wick, Foliar Wiper

Note: Best used when temps are below 85 degrees F, Rain-fast in two hours

Triclopyr Ester (Trade Name: Tahoe 4e)

Category: Selective

Target Species: Woody Plants – Upland / Terrestrial Areas

Application Methods: Basal Bark & Cut Stem

Note: Used year-round, except when temperatures are below 19 degrees F or when precipitation is taking place

Clethodim (Trade Name: Intensity)

Category: Selective

Target Species: Grasses – Upland / Terrestrial Areas

Application Methods: Foliar Broadcast Spray

Note: Used primarily during the cool seasons when native warm season grasses are dormant or not actively growing



Chemical Cocktail



Only the finest will do.





Stewardship



Advanced ID



Advanced ID



Work Commenced Fall 2011

Amos Tversky and Daniel Kahneman worked together to develop prospect theory, which aims to explain irrational human economic choices.

Richard Thaler – behavioral finance & mental accounting



10/06/2011 18:36

Thank you!

Doug Bauer
(847)533-5517
doughb@pizzo.info